

*CARELIA PILSBRYI*, N.SP., FROM THE HAWAIIAN ISLANDS.

By E. R. SYKES, B.A., F.L.S.

*Read 11th December, 1908.*

SHELL elongate, spire much produced, imperforate; sculpture consisting of well-marked, irregular, longitudinal lines, these gradually becoming weaker towards the protoconch, which is smooth; colour of the last whorl red-brown, gradually fading on the upper whorls till the protoconch is only faintly tinged. Whorls 8, the last measuring more than half the entire length, moderately inflated, with a well-marked suture; mouth lunate, outer lip (broken) hardly thickened at all, the white columellar margin reflexed and slightly expanded. Long. 65, lat. 23 mm.

*Hab.*—Kauai.

A single specimen, purchased from the collection of the late Mr. Rogers, has remained for some years unnamed in my collection. As other workers have also been unable to identify it, I now give a description and figure.

From *C. bicolor*, Jay, it differs in its larger size and also its colour; from *C. cochlea*, Rve., and *Cumingiana*, Pfr., in the absence of keeling and the more inflated whorls; from *C. Dolei*, Ancey, its size will readily distinguish it; from *C. olivacea*, Pease, it differs in colour and shape; from *C. paradoxa*, Pfr., it may be known by the absence of granular sculpture; from *C. turricula*, Mighels, it may be separated by the colouring, length of the last whorl, and lack of keel at the base. *C. Sinclairi*, Ancey, is unknown to me, and *C. glutinosa*, Ancey, is not, I believe, really a Hawaiian shell. I have only seen a single specimen of *C. Pilsbryi*, and the species is, I should think, an extinct one, like some others of the group.